

## ORIGINAL RESEARCH

**PHYTOCHEMICAL INVESTIGATIONS AND ANTIBACTERIAL AND ANTIOXIDANT PROPERTIES OF *THYMELAEA MICROPHYLLA* ESSENTIAL OIL****Souhila Bounab<sup>1</sup>, Takia Lograda<sup>1</sup>, Messaoud Ramdani<sup>1</sup>, Pierre Chalard<sup>2</sup> and Gilles Figueredo<sup>3</sup>**<sup>1</sup> *Laboratory of Natural Resource Valorisation, SNV Faculty, Setif 1 University, 19000 Setif, Algeria*<sup>2</sup> *SIGMA Clermont, Campus Des Cezeaux, CS 20265, 63178 Aubière Cedex, France*<sup>3</sup> *LEXVA Analytique, 460 Rue Du Montant, 63110 Beaumont, France***Submitted on: 10.12.17; Revised on: 15.12.17; Accepted on: 30.12.17****ABSTRACT**

The chemical composition of essential oil, isolated from *Thymelaea microphylla* by hydrodistillation, was analysed by GC and GC/MS. A total 30 compounds representing 99.91% of the oil were identified in Dreaat population. The essential oil of *T. microphylla* is characterized by a high rate of tridecanal (31.24%), nonanal <n> (11.43%), pentadecan 2-one (6Z) (7.93%) and the citronellol (6.80%). Antioxidant capacity was determined using the method of 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical. The extracts essential oil showed a good antioxidant activity. Disc diffusion method was used to evaluate the antibacterial activity on five different strains. The oil showed a significant effect against Gram-negative and positive bacteria.

**KEYWORDS:** Essential oil, Chemical composition, Antioxydant activity, Antibacterial activity, *Thymelaea microphylla*, Algeria

**Corresponding author:** Messaoud Ramdani  
**E-mail:** [ramdanimesaoud@yahoo.com](mailto:ramdanimesaoud@yahoo.com)  
**Tel:** +213 658 101010

**Indian Research Journal of Pharmacy and Science; 15(2017)1205-1215;**  
**Journal Home Page: <https://www.irjps.in>**  
**DOI: 10.21276/irjps.2017.4.4.7**

## INTRODUCTION

Most of the natural products reported and listed in the genus *Thymelaea*, belong to the large group of polyphenolic compounds and more precisely to flavonoids<sup>1-6</sup>. Compounds of the sterol family are reported in *T. microphylla*<sup>5</sup> and in *T. hirsuta*<sup>1</sup>. *T. microphylla*, *T. hirsuta*, *T. lythroides* and *T. tartonraira* are very rich in coumarines<sup>1,5,7</sup>. The microphybenzimidazole compound was isolated from *T. microphylla*<sup>8</sup>.

The essential oils of the genus *Thymelaea*, have been the subject of several phytochemical studies. *T. hirsuta* oils are characterized by high amounts of sesquiterpenes and monoterpenes<sup>9-10</sup>. The *T. microphylla* essential oil composition is dominated by the monoterpenes (67.84 %) <sup>11</sup>, and the study of its ethanolic extracts showed the presence of monoterpenes glycosides<sup>12</sup>.

*Thymelaea hirsuta* is known for its powerful antiseptic, hypoglycemic and anti-inflammatory properties and for the treatment of hypertension, such as antimelanogenesis, antidiuretic and purgative and in the treatment of influenza<sup>1-2,5-6,10,13-17</sup>. In M'Sila region, *T. hirsuta* is recommended by herbalists for the treatment of human diseases (Leishmanicide, dewormer and eczema)<sup>18</sup>. In Algeria anti-cancer and anti-inflammatory effects of *T. microphylla* methanol

extracts have been tested<sup>5,19-20</sup>. *Thymelaea lythroides* extracts have strong antifungal activity<sup>2</sup>.

*T. microphylla* essential oil showed a high antibacterial activity against *E. coli*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, and a weak inhibitory effect against *Pseudomonas aeruginosa*<sup>21</sup>. The essential oil of *T. hirsuta*, shows antioxidant, antimicrobial and antifungal activities<sup>9,16,19</sup>. The cytotoxicity of the oil of this species is demonstrated by Felhi *et al.*<sup>16</sup>.

The objective of this study is to determine the chemical composition of *Thymelaea microphylla*, species endemic to the Sahara, and to evaluate the antibacterial and antioxidant activities of its essential oil.

## MATERIALS AND METHODS

### Plant material


*Thymelaea microphylla* Coss. & Dur. commonly called "Methnane Ghazal or Methnane Labiadh", is a plant endemic to North Africa. It is a woody plant with much branched stems whose leaves are small, ovoid, scattered and distant on the branches. Flowers are glomerulate 2-5 in the axils of leaves, yellowish in color with very short lobes. This plant is found in arid and desert pastures<sup>22</sup> (Figure-1).



Figure 1: *Thymelaea microphylla* from the M'Sila region.

*Thymelaea microphylla* is collected from the Dreat forest (Hammam Dalaa, M'Sila). Aerial parts were

collected during the flowering stage in Mai 2017 (Figure 2).



**Figure 2:** Sampling area of *T. microphylla* Coss. and Dur.

The air dried materials were subjected to hydro-distillation for 3h using a Clevenger apparatus type. Voucher specimens were deposited in the herbarium of the Department of Biology and Ecology, Setif University, Algeria. The oil obtained was collected and dried over anhydrous sodium sulphate and stored in screw capped glass vials in a refrigerator at 4-5°C prior to analysis. Yield based on dried weight of the samples was calculated.

#### Essential oil analysis

The essential oils were analysed on a Hewlett-Packard gas chromatograph CPG/FID 7890, coupled to a gaz chromatograph: CPG/MS 7890/5975C, equipped with a Colonn Apolar: DB5 MS: 40 m 0,18 mm 0,18 µm, programming from 50 °C for 5min – 5 °C/min until 300 °C. Helium was used as the carrier gas (1.0 ml/min); injection in split mode (1:30), injector and detector temperature is 280 °C with split 1/100. The mass spectrometer worked in EI mode at 70 eV; electron multiplier, 2500 V; ion source temperature, 180°C; MS data were acquired in the scan mode in the *m/z* range 33-450. The identification of the components was

based on comparison of their mass spectra with those of NIST mass spectral library<sup>23-24</sup> and those described by Adams as well as on comparison of their retention indices either with those of authentic compounds or with literature values<sup>25</sup>.

#### Antibacterial Activity

The Extract Essential oil was tested against the following bacteria; 2 gram negative bacteria: *Escherichia coli* ATCC 25922; and *Pseudomonas aeruginosa* ATCC 27853, and 3 gram positive bacteria; *Bacillus subtilis* ATCC 6633, *Staphylococcus aureus* ATCC 25923 and *Enterococcus faecalis* ATCC 51299. The in vitro antibacterial activity of the examined extract was assessed the determination of the activity by the micro dilution method, according to recommendations of the Clinical and Laboratory Standards Institute (NCCLS). The bacterial inoculum was prepared from overnight broth culture in physiological saline (0.8 % of NaCl) in order to obtain an optical density ranging from 0.08-01 at 625 nm. Muller-Hinton agar (MH agar)

and MH agar supplemented with 5 % sheep blood for fastidious bacteria were poured in Petri dishes, solidified and surface dried before inoculation. Sterile discs (6 mm ) were placed on inoculated agars, by test bacteria, filled with 10 l of mother solution and diluted essential oil (1:1, 1:2, 1:4, 1:8 and 1:16 v:v of DMSO). DMSO was used as negative control. Bacterial growth inhibition was determined as the diameter of the inhibition zones around the discs. All tests were performed in triplicate. Then, Petri dishes were incubated at 37°C during 18 to 24h aerobically (Bacteria). After incubation, inhibition zone diameters were measured and documented.

#### Antioxidant activity

The free radical-scavenging activity of *T. microphylla* crude extract was measured in terms of hydrogen donating or radical-scavenging ability using the stable radical DPPH. In this assay, the purple chromogen radical 2,2-diphenyl-1-picrylhydrazyl (DPPH) is reduced by antioxidant/reducing compounds to the corresponding pale yellow hydrazine<sup>26</sup>. The scavenging capacity is generally evaluated in organic media by monitoring the absorbance decreases at 517nm until the absorbance remains constant. 4.0 mM solution of DPPH in methanol was prepared and 1.0 ml of this solution was added to 2.9 ml of extract solution in methanol at different concentrations (37.5 to 300 µl). Thirty minutes later, the absorbance was measured at 517 nm. BHT (Butelated hydroxytoluen) was used as the standard. Lower absorbance of the reaction mixture indicated higher free radical scavenging activity. Radical-scavenging activity was expressed as the

inhibition percentage of free radical by the sample and was calculated using the following formula<sup>27</sup>:

$$I\% = \left( \frac{A.blank - A.sample}{A.blank} \right) * 100$$

Where (A.blank) was the absorbance of the control (blank, without essential oil) and (A.sample) was the absorbance in the presence of the Essential oil. All the tests were performed in triplicate and the graph was plotted with the mean values.

## RESULTS

### 1. Chemical analysis

The hydro-distillation of *Thymelaea microphylla* essential oil gave a viscous liquid with pale yellow oil. The average yield of essential oil of the sample is 0.14%. The analysis and identification of the components of the essential oil of this species was performed using the GC-GC/MS (Figure 3). The compound identified in these oils and the abundances are presented in order of their appearance (Table 1).

This analysis led to the identification of 30 components representing 99.93% of the total oil of *T. microphylla*. According to our results the chemical composition of the species, *T. microphylla* is dominated by the presence of major compounds, the tridecanal (31.24%), nonanal<n-> (11.43%), pentadecen 2-one (6Z) (7.93%) and the citronellol (6.80%). As well as other component with lower percentages, tetradecanal (4.43%), caryophyllene 14, 6-hydroxy 4-5 dihydro (3.65%), lavandulyl acetate tetrahydro (3.50%), dodecanal (3.27%), undecanal (3.09%) and the presence in trace of other compounds.




Figure 3: GC/Masse and GC/FID profiles of *T. microphylla*



**Table 1:** Chemical composition of *Thymelaea microphylla* essential oil

Yield %	KI	0.14	Yield %	KI	0.14
Number of compound		30	Number of compound		30
Total		99.91	Total		99.91
Pentanone-4-hydroxy-4-methyl-2	838	1.76	Citronellyl acetate	1322	0.89
$\alpha$ -pinene	935	0.55	Tetradecane (C14)	1399	1.55
Pentyl furan-2	989	0.62	Dodecanal	1410	3.27
Octanal	1002	0.58	Neryl acetone	1449	2.27
Nonanal-n	1102	11.43	Dodecen 1-ol (2E)	1476	1.41
Camphor	1151	0.56	$\beta$ -ionone (E)	1483	1.11
Lavandulol	1164	0.57	Tridecanal	1513	31.24
Naphthalene	1191	1.94	Hexenyl benzoate (3Z)	1577	0.54
Dodecane	1200	2.80	Tridecen 1-al (2E)	1585	1.74
Decanal-n	1207	1.64	Sesquisabinene hydrate Trans	1595	0.57
Citronellol	1224	6.80	Tetradecanal	1614	4.43
Lavandulyl acetate tetrahydro	1273	3.50	Isomenthone 2(3-oxobutyl)	1679	1.35
Isopulrgyl acetate iso	1295	1.06	Pentadecen 2one (6Z)	1716	7.93
Tridecane	1300	0.55	Caryophyllene 14, 6 hydroxy 4-5 dihydro	1841	3.65
Undecanal	1309	3.09	Farnesyl acetone (5E, 9Z)	1912	0.51

**2- Antibacterial activity**

The results of the experiments assessing the bacteriostatic effects of *T. microphylla* essential oil

of the study on Gram-negative and Gram-positive bacteria are presented in (Table 2).

**Table 2:** Antibacterial activity of *T. microphylla* essential oil

Bacteria	Antibiotic*		Dilution				
	Gen	Amo	1/1	1/2	1/4	1/8	1/16
<i>Bacillus subtilis</i> ATCC 6633	26	-	29	19.5	18.5	14	11.5
<i>Escherichia coli</i> ATCC 25922	26	-	26.5	26	16.5	15.5	13.5
<i>Enterococcus faecalis</i> ATCC 51299	-	26	18.5	17.5	15.5	15	14.5
<i>Pseudomonas aeruginosa</i> ATCC 27853	28	-	-	-	-	-	-
<i>Staphylococcus aureus</i> ATCC 25923	30	-	-	-	-	-	-

\* Gen= Gentamicine; Amo= Amoxicilline

Effectively, the essential oil from *T. microphylla* aerial part were demonstrated antibacterial activity against the gram negative clinical pathogens bacteria tested (Figure 4),

*Bacillus subtilis*, *Escherichia coli* and *Enterococcus faecalis*, but no activity against the bacteria *Pseudomonas aeruginosa* and *Staphylococcus aureus*




Figure 4: Antibacterial activity of *T. microphylla* essential oil

### 3- Antioxidant activity

The antiradical activity of *T. microphylla* essential oil is evaluated by their inhibitory capacity of a methanolic solution of DPPH; it's measured by spectrophotometer at 517 nm. The standard used

was BHT (Butylated hydroxytoluen) (Table 3). The change in the color of the solution containing *T. microphylla* essential oil proves that the oil has antioxidant activity.

Table 3: Antioxidant activity of essential oil of *T. microphylla*

Concentrations	300 µl (1/1)	150 µl (1/2)	75 µl (1/4)	37.5 µl (1/8)
Absorbance of EO (nm)	0.057	0.119	0.317	0.515
Absorbance de BHT (nm)	0.068	0.14	0.330	0.640
Inhibition (%) of EO	91.53	82.31	52.89	23.47
Inhibition (%) of BHT	89.89	79.19	50.96	4.90

The inhibition percentages of different concentrations of *T. microphylla* essential oil as well as the BHT standard fluctuate between 4.90% and 89.89% for BHT, 23.47% and 91.53% for *T.*

*microphylla* oil (Figure 5). It should be noted that there is a certain similarity in the percentages of inhibition, for the oil and for the BHT standard, except for the 1/8 dilution.




Figure 5: Inhibition percentage of *T. microphylla* essential oil

The highest inhibition value is 91.53% for oil and 89.89% for BHT, indicating a possibility that *T. microphylla* oil contains a greater amount of free radical accepting compounds, as well as that the greatest antioxidant potential. The percentage of inhibition of the free radical increases with the increase of the concentration, either for the BHT or for the essential oil. IC50 is inversely related to the antioxidant capacity of a compound because it expresses the amount of antioxidant required to decrease the free radical concentration by 50%.

#### DISCUSSION :

Many works have shown a great heterogeneity in the chemical composition of the genus *Thymelaea*. In Tunisia, *T. hirsuta* is characterized by heptane, germacrene-D,  $\gamma$ -eudesmol and citronellyl formate<sup>9,28</sup>. The *T. hirsuta* essential oil, cultivated in Tunisia, is dominated by hexadecanoic acid, 4, 8- and 13- dimethylhecosane methylhexacosane<sup>10</sup>.

Chemical analysis of *T. microphylla* essential oils allowed us to identify tridecanal (31.24%), nonanal <n-> (11.43%), pentadecan (7.93%) and citronellol (6.80%) as major components. The chemical results differ from those cited in the literature. *T. microphylla* samples from the Ourgla region (Algeria) show the predominance of D-menthone, 2-undecanone, pulegone and perillal<sup>11</sup>. This difference in composition is probably due to various conditions including the environment, geographical origin, harvest period, temperature and drying time.

*T. hirsuta* essential oil shows significant antioxidant activity<sup>17,29</sup>. This oil reduces the formation of DPPH radicals in a dose-dependent manner<sup>9</sup>. The investigation of the antioxidant activity of the essential oil of *T. microphylla* has shown that this species has antioxidant activity. The various studies of aqueous and methanolic extracts

of *T. microphylla* prove the presence of antioxidant activity<sup>3,6,12,20-21,30</sup>.

Species of the genus *Thymelaea* exhibit important biological activities. The essential oil of *T. hirsuta* has moderate antimicrobial activity against all microorganisms tested<sup>16,19</sup>. *T. lythroides* has a strong antifungal activity<sup>2</sup> where as the methanolic extracts of *T. microphylla* showed no bacteriological effect<sup>20</sup>.

Our investigation shows a high activity of *T. microphylla* essential oil on *Bacillus subtilus*, *Echerichia coli*, *Enterococcus faecalis* bacteria and no effect against *Pseudomonas aeruginosa* and *Staphylococcus aureus*.

The studies of Noman *et al.*<sup>31</sup> showed high antibacterial activity of *T. microphylla* essential oil against *E. coli* and *S. aureus*, *K. pneumoniae* and a weak inhibitory effect against *P. aeruginosa*. The absence of antibacterial activity can be explained by the developed resistance of some strains that react differently to several essential oils; this is the case of *P. aeruginosa*<sup>32-33</sup>.

#### CONCLUSION :

This study aims to identify the chemical composition of *Thymelaea microphylla* essential oils and the evaluation of their antioxidant and antibacterial activities. The analysis of the chemical composition of the essential oil by GC/MS has allowed the identification of 30 compounds. The tridecanal is the major component of the chemical composition, although the study has identified the other natural products as minor components. The results exhibited antioxidant activity, which are stronger than the positive control BHT. An additional characteristic of *T. microphylla* essential oil was its prominent antibacterial activity against *B. subtilus*, *E. coli*, *E. faecalis* and no effect against *P. aeruginosa* and *S. aureus*. The Essential



oil may be a promising alternative treatment of localized infections even with severe hospital acquired strains. Further investigations must be made to determine the active constituent(s) for their application in medical research.

#### REFERENCES:

1. Dohou N, Yamni K, Tahrouch S, Idrissi Hassani LM, Badoc A et Gmira N., Screening phytochimique d'une endémique ibéro-marocaine, *Thymelaea lythroides*. Bull. Soc. Pharm. Bordeaux, **2003**; 142: 61-78.
2. Dohou N, YAMNI K, Badoc A, Douira A., Activité antifongique d'extraits de *Thymelaea lythroides* sur trois champignons pathogènes du RIZ. Bull. Soc. Pharm. Bordeaux, **2004** ; 143: 31-38.
3. Benhammou N, Atik Bekkara F and Jean-Marie Coustard, Extracts from antioxidant activity of methanolic and water extracts from *Marrubium deserti* (de Noë) and *Thymelaea microphylla* from Algerian Sahara. Advances in Food Sciences, **2009**; 31(4): 194-201.
4. Ghanem H, Haba H, Marcourt L, Benkhaled M and Wolfender J-L., Microphynolides A and B, new spiro- $\gamma$ -lactone glycosides from *Thymelaea microphylla* Coss et Dur., Nat Prod Res., **2014**; 28(20):1732-1738.
5. Mekhelfi T., Séparation et Détermination Structurale de Métabolites Secondaires de deux Plantes Algériennes - Activités Biologiques. Thèse de Doctorat en Chimie pharmaceutique. Université de Constantine, Algérie, **2016**; 220p.
6. Kerbab K., Validation scientifique de l'utilisation traditionnelle des plantes médicinales algériennes: *Halimium halimifolium* (L.)Wilk et *Thymelaea microphylla* Coss.et Dur.,

#### ACKNOWLEDGEMENTS:

The work was supported by Algerian MESRS and Chemical Laboratory of carbohydrates Heterocyclic of Clermont Ferrand, France.

- composition chimique et activité biologique. These de Doctorat en Chimie pharmaceutique. Université des frères Mentouri-Constantine, **2017**; 239p.
7. Mekhelfi T, Kerbab K, Guella G, Zaiter L, Benayache S, and Benayache F., Phytochemical constituents of *Thymelaea microphylla* Coss. et Dur. from Algeria Der Pharmacia Lettre, **2014**; 6(1): 152-156.
  8. Noman L, Oke-Altuntas, Zellagui A, Ayse Sahin Yaglioglu, Demirtas I, Cardoso S M, Akkal S, Gherraf N et Rhouati S., A novel benzimidazole and other constituents with antiproliferative and antioxidant properties from *Thymelaea microphylla* Coss. et Dur. Natural Product Research. Formerly Natural Product Letters, **2017**; 31(17): 2032-2041.
  9. Kadri A, Zarai Z, Ben Chobba I, Gharsallah N, Damak M et Békir A., Chemical composition and in vitro antioxidant activities of *Thymelaea hirsuta* L. essential oil from Tunisia, African Journal of Biotechnology, **2011**; 10(15): 2930-2935.
  10. Yahyaoui M, Bouajila J, Camy S, Condoret JS and Abderabba M., Comparaison of chemical composition and yield of *Thymelaea hirsuta* volatile oils obtained by supercritical carbon dioxide extraction and hydrodistillation methods. International Symposium on Essential oils natural volatiles & Essential oils, Istanbul, **2014**; Turkey.

11. Noamane SL, Zellagui A, Mesbah K, Gherraf N, Lahouel M and Rhouati S., Essential oil Composition of *Thymelea microphylla* Coss et Dur Der Pharmacia Lettre, **2010**; 2(5): 428-431.
12. Kerbab Khawla, Tarak Mekhelfi, Lahcene Zaiter, Samir Benayache, Fadila Benayache, Patrizia Picerno, Mencherini T, Sansone F, Aquino R.P et Rastrelli L., Chemical composition and antioxidant activity of a polar extract of *Thymelaea microphylla* Coss. & Dur., Natural Product Research (Formerly Natural Product Letters), **2014**; 29(7): 671-675.
13. Ziyat A, Legssyer A, Mekhfi H, Dassouli A, Serhrouchni M, Benjelloun W., Phytotherapy of hypertension and diabetes in oriental Morocco, J. Ethnopharmacol., **1997**; 58: 45-54.
14. Dohou N, Yamni K, Gmira N et Lalla Mina Idrissi Hassani, Étude de Polyphénols des feuilles d'une endémique ibéro-marocaine: *Thymelaea lythroides*, Acta Botanica Malacitana., **2004**; 29: 233-239.
15. Bnouham M, Benalia W, Bellahcen S, Hakkou Z, Ziyat A, Mekhfi H, Aziz M and Legssyer A., Antidiabetic and antihypertensive effect of a polyphenolrich fraction of *Thymelaea hirsuta* L. in a model of neonatal streptozotocin-diabetic and NG-nitro-l-arginine methyl ester-hypertensive., Journal of Diabetes., **2012**; 4: 307-313.
16. Felhi S, Chaaibia M, Bakari S, Ben Mansour R, Békir A, Gharsallah A and Adel Kadri A., Anti-microbial screening and cytotoxic activity of aerial part of *Thymelaea hirsuta* L. essential oil growing in south-west Tunisia. Pak. J. Pharm. Sci., **2017**; 30(1): 087-091.
17. Yahyaoui M, Ghazouani N, Ines Sifaoui I, and Manef Abderrabba, Comparison of the Effect of Various Extraction Methods on the Phytochemical Composition and Antioxidant Activity of *Thymelaea hirsuta* L. aerial parts in Tunisia. Biosciences biotechnology research Asia, **2017**; 14(3): 997-1007.
18. Boudjelal A, Henchiri C, Sari M, Sarri D, Hendel N, Benkhaled A and Ruberto G., Herbalists and wild medicinal plants in M'Sila (North Algeria): An ethnopharmacology survey. Journal Ethnopharmacol., **2013**; 148(2): 395-402.
19. Dahamna S, Dehimi K, Merghem M, Djarmouni M, Bouamra D, Harzallah D, Khennouf S., Antioxidant, Antibacterial and Hypoglycemic Activity of Extracts from *Thymelaea microphylla* Coss. et Dur. International Journal of Phytocosmetics and Natural Ingredients. International Journal of Phytocosmetics and Natural Ingredients, **2015**; 2: 15-18.
20. Dehimi K, Antonio Speciale, Antonina Saija, Saliha Dahamna, Roberto Raciti, Francesco Cimino, Mariateresa Cristani. 2016. Antioxidant and anti-inflammatory properties of Algerian *Thymelaea microphylla* Coss and Dur. Extracts. Pharmacognosy Magazine., **2016**; 12(47): 203-210.
21. Noman L, Amar Zellagui A, Youcef Hallis Y, Sahin Yaglioglu, Ibrahim Demirtas I, Nouredine Gherraf N and Rhouati S., Antioxidant and antimicrobial activities of an endemic desert species *Thymelea microphylla* Coss. et Dur. Der Pharmacia Lettre., **2015**; 7(1): 118-121.
22. Quézel P et Santa S., Nouvelle Flore de l'Algérie et des régions désertiques méridionales, Ed. CNRS, Paris, 2 Vol. **1963**.

23. Masada Y., Analysis of essential oils by Gas Chromatography and Mass Spectrometry. *J. Wiley & Sons, Inc. New York*, **1976**.
24. NIST., Mass Spectral Search Program for the NIST/EPA/NIH Mass Spectral Library, vers. 2.0. fiveash data, USA, **2002**.
25. Adams RP., Identification of essential oils components by Gas Chromatography-Mass spectroscopy, Allured Publishing Corporation Carol Stream, Illinois USA, **2007**.
26. Epifano F, Genovese S, Menghini L and Curini M., Chemistry and pharmacology of oxyprenylated secondary plant metabolites. *Phytochemistry*, **2007**; 68: 939-953.
27. Archana B, Dasgupta N and De B., In vitro study of antioxidant activity of *Syzygium cumini* fruit. *Food Chem.*, **2005**; 90: 727-733.
28. Ben Chobba I, Felhi S, Chaaibia M, Ben Mansour R, Békir A, Drira N, Gharsallah N and Kadri A., Phytochemical composition and biological assays of the aerial parts of *Thymelaea hirsuta* essential oil from Tunisia. The Third International Symposium on the Biology of Rare and Endemic Plant Species (BIORARE-2014) April 19-23, **2014**; Antalya, Turkey.
29. Amari ON, Bouzouina M, Berkani A and Lotmani B., Phytochemical screening and antioxidant capacity of the aerial parts of *Thymelaea hirsuta* L. *Asian Pac. J. Trop Dis. Apr.*, **2014**; 4(2): 104-109.
30. Benhammou N., Activité antioxydante des extraits des composés phénoliques de dix plantes médicinales de l'Ouest et du Sud-Ouest Algérien. Thèse de Doctorat en Biologie. Option: Substances Naturelles, Activités biologiques et Synthèse. Université Aboubakr Belkaïd-Tlemcen (Algérie), **2012**; 174 p.
31. Noman L, Zellagui A, Ayse Sahin Yaglioglu., Demirtas I and Rhouati S., Antimicrobial Activity of Essential Oil Components and Antiproliferative Activity of Trans-Tiliroside Compound from an Endemic Desert Species *Thymelaea microphylla* Coss. et Dur. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, **2015**; (2): 671-676.
32. Mann CM, Cox SD and Markham JL., The outer membrane of *Pseudomonas aeruginosa* NCTC 6749 contributors to its tolerance to the essential oil of *Melaleuca alternifolia* (Tea tree oil). *Letters in Applied Microbiology*, **2000**; 30: 294-297.
33. Hammer KA, Carson CF and Riley TV., Antimicrobial activity of essential oils and other plant extracts, *Journal of Applied Microbiology*, **1999**; 86: 985-990.

CONFLICT OF INTEREST REPORTED: NIL ;

SOURCE OF FUNDING: NONE REPORTED