

Original Research



ASSESSMENT OF KNOWLEDGE, ATTITUDE AND PERCEPTIONS REGARDING PHARMACOVIGILANCE AMONG THIRD TERM MEDICAL STUDENTS IN SOUTHERN INDIA

Harish. G. Bagewadi^{*1}, Venkatadri.T.V²

^{*1}Asst. Professor, ²Professor & HOD- Department of Pharmacology, MVJ Medical College & Research Hospital, Bangalore, India -562114

Submitted on: 25.05.2015

Revised On: 30.05.2015

Accepted on: 01.06.2015

ABSTRACT:

Aim- To assess the awareness of Pharmacovigilance, Adverse Drug Reaction (ADR) reporting, and to evaluate the impact of an educational intervention among third term medical students.

Materials and methods- This was a questionnaire-based, pre- and post-test educational interventional study. All the students were given handouts containing information about pharmacovigilance and ADR reporting one week before the educational intervention. A pre-validated 20-point questionnaire on (KAP) Knowledge, attitude, perception about Pharmacovigilance and Adverse Drug Reactions (ADR's) reporting was distributed to third term medical students (n=52). An interactive educational intervention using overhead projector and transparencies (OHP) was designed. The chi-square test and unpaired paired t-test was used for statistical calculation.

Results- The overall response rates were expressed as percentages, Mean±SD. The knowledge, attitude and perceptions of pharmacovigilance and adverse drug reactions when compared before (pre-KAP) and after (post-KAP) the educational intervention, the correct response rates were found to be statistically significant (P<0.001). The feedback from the students was encouraging and positive. The handouts before the lecture classes helped them to easily grasp the pharmacovigilance and adverse drug reactions concepts better during lectures.

Conclusion- The study concluded that imparting the knowledge about pharmacovigilance and ADR reporting can play vital role in drug safety and rational use of medicines in future.

KEYWORDS: KAP questionnaire, Pharmacovigilance, ADR reporting, Continuous medical education.

Corresponding Author- Harish G Bagewadi

Email: harish2957@gmail.com

Telephone: +91 8884376427

Indian Research Journal of Pharmacy and Science; 5(2015) 180-186;
Journal home page: <https://www.irjps.in>

INTRODUCTION:

The safe and cost effective medicines are high requisitions in the modern world. In 1968, the first practical international co-operation in drug monitoring was established. The ideas came up as a consequence of the so called thalidomide tragedy. In the 1960's it was discovered that if thalidomide is ingested by mothers during pregnancy limb deformities in babies may occur. This incident became the modern starting point of a science focusing on patient problems due to medicinal use. Medication safety is a more significant issue, because of immense competition among pharmaceutical manufacturers; medicinal products may be registered and marketed in many countries simultaneously. As a result, adverse reactions may not always be readily identified and so are not monitored systematically. Pharmacovigilance has constantly grown its importance in last 15 years, relating to the absolute amount of adverse drug reactions (ADR's) and to the fact of several hospital admissions are due to ADRs^{1, 2}. Pharmacovigilance is an arm of patient care and surveillance. It aims at getting the best outcome from treatment with medicine. Adverse drug reactions (ADRs) are common causes of morbidity and mortality in both hospital and community settings. Adverse drug reactions (ADRs) are global problems of major concern. ADRs are responsible for about 5% to 20% of hospital admissions^{3,4}. World Health Organization (WHO) defines ADR as "any response to a drug which is noxious and unintended, and which occurs at doses normally used in man for prophylaxis, diagnosis, or therapy of disease, or for the modification of physiological function".

Studies from different settings indicate inadequate knowledge about pharmacovigilance among healthcare professionals as well as attitude that are associated with high degree of underreporting⁵⁻¹⁰. Assessment of awareness of Pharmacovigilance among the healthcare professionals is very important due to under reporting of adverse drug reactions. Ensuring the safe use of drugs is a combined responsibility of the healthcare team that includes doctors, nurses, pharmacists and other supporting staff¹¹. As future medical practitioners, medical students need to be well trained on how to recognize, prevent and report ADRs. Therefore, the aim and objective of this study was to evaluate knowledge, attitude and the perceptions about Pharmacovigilance among medical students at medical college in south India by an interactive educational module as an intervention.

MATERIALS AND METHODS:

The study was conducted at MVJ medical college and research hospital, Bangalore. Permission was

duly taken from Institutional Ethics Committee to conduct the study. This was a prospective, Knowledge, Attitude and Perceptions (KAP) questionnaire based study. Third term medical students (n=52), were participated in the study. The students were given handouts containing information about pharmacovigilance and ADR reporting one week before the educational intervention, which was designed using overhead projector and transparencies (OHP) for one hour by trained faculty to all participants. The Pre-KAP questionnaire survey was done before educational intervention in order to facilitate the transfer of knowledge of pharmacovigilance and ADR's reporting. The educational intervention consisted of a theoretical content about what is pharmacovigilance, its main objectives, adverse drug reactions reporting, vigiflow database, classification of ADRs, incidence of ADRs, role of health care professionals, reporting of suspected adverse drug reaction followed by economic and epidemiological importance of reporting the ADRs and its effect on patient safety and causality assessment of ADRs.

A Semi-structured, pre-validated, questionnaire was used for data collection as a research tool. A structured questionnaire was designed after minor modifications from the work of V. Lokesh Reddy et al¹², Radhakrishnan Rajesh et al¹³ and Harish .G. Bagewadi et al¹⁴. The KAP questionnaire (Annexure - 01) consisted of 20 questions about pharmacovigilance and ADR's reporting. Out of which, 11 questions were related to knowledge, 05 questions were related to attitude, 04 questions were related to perception. The correct responses were scored 1 point and wrong responses were given zero point for knowledge related questions and practice related questions. The attitude related questions were scored based upon the participant's degree of agreement using Likert scale. The score was as following; "0"- strongly disagree, "1" - disagree, "2"- uncertain, "3"- agree and "4"- strongly agree. In order to preclude any potential bias, the disclosure of name of the responder was made optional.

Before the start of educational intervention, initially all the students were briefed about the purpose of the study, student's consent was taken; later pre-KAP questionnaire was administered and asked to submit the same. After the interactive educational intervention program on pharmacovigilance, all participants of Pre-KAP questionnaire in the study was administered with Post-KAP questionnaire and it was analyzed, question wise and their responses were documented.

The filled KAP questionnaires were evaluated as per the study objectives, the KAP scores were analyzed. The data obtained were entered in Microsoft excel spread sheet and evaluated. The impact of effectiveness of educational intervention on the awareness of pharmacovigilance and ADR's reporting among the third term medical students is evaluated. The chi-square test and unpaired t test was used to compare the difference in correctness for each question. All results attained were entered in Microsoft excel and the statistical calculations were executed using Graph Pad InStat. The p value ($p < 0.05$) is considered to be statistically significant.

RESULTS:

All the answers are expressed in terms of numbers, percentages and Mean \pm SD, for the KAP questionnaire (Pre-KAP & Post-KAP) comprising of 20 questions was evaluated and tabulated in Table.1, 2, 3, 4 and Figure 1.

Question 01 of table 01, emphasized on the role of health care professionals in ADR's reporting, for which the comparativeness with educational intervention in between correct response score of pre-KAP (33.3%) and post-KAP (74.1%) revealed effective educational intervention with statistically significance ($p < 0.0001$). In our study it was demonstrated by an increase in the correct responses for pre and post KAP question 02, table 01 in defining pharmacovigilance, from 37.1% before to 66.6% after the intervention. Question 03 from table 01 framed to obtain the information about objectives of pharmacovigilance where in pre-KAP (20.4%) and post-KAP (63%) and statistically significant ($p < 0.0001$) correct responses were evident after educational programme. This data suggests that continuing educational intervention is an important tool for increasing all health care professionals' awareness to pharmacovigilance. Based on our study results and the finding of Cosentino et al¹⁵ and Figueras et al¹⁶ recommend including "pharmacovigilance" as a topic in continuing education programmes and would also recommend a yearly repetition of such educational interventional program to all health care professionals.

Question 04 from table 01 is framed to obtain the information about the international center for reporting of ADR's and question 06 from table 01, about databases on ADR's reporting system, it was found that there was an increased positive response rate of 18.5% as pre-KAP to 63% post-KAP and 13% as pre-KAP to 48.15% post-KAP after the educational intervention program respectively. The result strongly suggests that students were greatly influenced by the educational intervention regarding the reporting systems of ADR's both of national and

international standards which is in accordance with earlier study by Suveges LG et al¹⁷.

Question 05 from table 01, is framed to obtain the causality assessment of an ADR, we could see drastic increased positive response of 16.6% before to 52.3% and after the educational intervention. Question 07 from table 01 is designed to know the information about rare ADR's found in clinical trials, we observed that increased positive response of 18.5% before to 59.26% after the educational intervention which points to create more awareness about phases of clinical trials and safety of medicines in medical students.

The study also focused on assessing the attitude of medical students- ADR's reporting in question 14 and 15 table no.2, which revealed to be 31.5% before pre-KAP to 92.6% post-KAP, 27.7% before pre-KAP to 85.2% post-KAP respectively, which strongly suggests that students need to undergo educational sessions on ADR's reporting¹⁸.

The study is focused on assessing the perception of ADR's reporting centers in question 17 table no.3, which revealed to be 27.7% before pre-KAP to 74.1% post-KAP, which also points to the importance of impact of educational interventions on Pharmacovigilance in accordance with earlier study by Scott HD et al¹⁸.

Question 18 table no 03, highlights on communication of safety information between all health care professionals which can minimize the risk of marketed medicines, observed to be 31.5% before pre-KAP to 77.7% post-KAP, students were made aware that communication among other health care professionals is important. Question 19 from table 03, showed that 33.3% before pre-KAP to 81.5% post-KAP depicts to change the perception of students. Because, rational use of medicines is the need of the hour by imparting knowledge about safety of drugs through educational interventions.

Question 20 from table 03, to read an article (online /newspaper/magazine) about ADR's in near future revealed to be 37% before pre-KAP to 70.4% post-KAP, this type of enthusiasm, motivation gained among students after educational intervention is noteworthy.

DISCUSSION:

The present study revealed that medical students who attended the interactive educational intervention on Pharmacovigilance and ADR's reporting were much satisfied, and considered it more effective and valuable. In our study, one focus of educational intervention was to increase the medical students awareness to Pharmacovigilance, regulatory bodies responsible for monitoring of ADR's, types of ADR's. This was demonstrated by an increase in the correct

responses in pre and post-KAP questions (1 to 20) about pharmacovigilance and ADR's reporting with statistical significance ($p < 0.0001$), after the educational intervention highlighting the impact on its effectiveness.

Questions 08 and 09 from table 01 were framed to obtain the knowledge about ADR and its causative drug which medical students, physicians must know to promote safe and rational use of medicines. The response rate is 27.7% as pre-KAP to 59.26% post-KAP and 33.3% as pre-KAP to 74.1% post-KAP respectively, after the educational intervention program. Question 10 from table 01, shows response rate from 37% pre-KAP to 70.4% post-KAP which strongly suggests that the information about different types of ADR, and question 11 from table 01, infers about when to report ADR's and practical knowledge on ADR from 40.7% pre-KAP to 64.8% post-KAP improved enormously after educational intervention. Question 12 from table 02, showed that 48.2% before pre-KAP to 77.7% post-KAP, and Question 13 from table 02 showed that 35.2% before pre-KAP to 79.6% post-KAP strongly suggests that there is a great need to create awareness on attitude aspect of ADR reporting among medical students can be done by continuous medical education programs on pharmacovigilance.

In figure 01, the total Pre-KAP scores on knowledge (2.8 ± 2.24), attitude (8.25 ± 5.18), perception (1.32 ± 1.75) when compared to total post-KAP scores on knowledge (7.4 ± 3.32), attitude (13.27 ± 3.24), perception (3.14 ± 4.26) respectively. The overall increase in correct response rate with statistical significance ($p < 0.0001$) was observed after educational intervention.

Earlier studies by Suveges LG et al and Scolt HD et al^{17,18} has also shown that enhancing knowledge,

attitude, and perception of improving awareness can increase the number of ADR reports. This study conducted by Chatterjee et al¹⁹ which stated that a main reason for under reporting of ADR's was the clinical negligibility of the adverse reaction due to lack of time and little knowledge about the types of reactions to be preferentially reported. However, in a similar educational interventional program in pharmacovigilance study of Li Q, Zhang et al²⁰ showed that educational intervention improved awareness of pharmacovigilance on knowledge, attitudes, practice of healthcare professionals.

The feedback from the students was encouraging and positive. The hand outs before the lecture classes helped them to understand the concepts better and potentiated easy grasping habits during lecture hours. Students are of the opinion that handouts when given before every Pharmacology lecture would help them to absorb concepts better during lecture classes. The photo images showing some examples of (ADR's and its causative drug) during lecture class, made students to learn ADR's causality effectively and to assess benefit/risk ratio of marketed medicines.

The study has some limitations. The study period was too short. The study findings could not be applied to the wider community medical students and other health care professionals as the study was restricted to third term medical students in department of Pharmacology, MVJ Medical College and Research hospital, Bangalore. Therefore we recommend that several such studies of similar kind should be conducted among wider community medical students as well as to all types of health care professionals so as to develop strategies to improve the knowledge, attitudes, practice of pharmacovigilance in India and globally.

Table.1. Knowledge of Pharmacovigilance & ADR reporting before & after educational intervention.

Q. no	K A P - Items	Pre-KAP Score n(%)	Post-KAP score n(%)	p-value
1.	The healthcare professionals responsible for reporting ADR in a hospital is/are- a) Doctor b) Pharmacist c) Nurses d) All of the above*	18 (33.3)	40 (74.1)	$p < 0.0001$
2.	Define Pharmacovigilance? a) The science of monitoring ADR's in the Hospital b) The process of improving the safety of Drugs c) The detection, assessment, understanding & prevention of adverse effects* d) The science of detecting the type & incidence of ADR after drug is marketed.	20 (37.1)	36 (66.6)	$p < 0.0001$
3.	The important objective of Pharmacovigilance is a) To identify safety of drugs* b) To calculate incidence of ADR's c) To identify predisposing factors to ADR's d) To identify ADR's occurring at high doses	11 (20.4)	34 (63)	$p < 0.0001$

4.	The international center for adverse drug reaction monitoring is located in: a) Unites States of America b)Australia c) Canada d) Sweden*	10 (18.5)	34 (63)	p<0.0001
5.	Which of the following scales is commonly used to assess the causality of an ADR? a)Hartwig scale b)Naranjo algorithm* c)Schumock & Thornton scale d)Karch & Lasagna scale	9 (16.6)	32 (52.3)	p<0.0001
6.	Which one of the following is the 'WHO online database' for reporting ADRs? a) ADR advisory committee b) Med safe c)Vigibase*d) Med watch	7 (13)	26 (48.15)	p<0.0001
7.	Rare ADRs can be identified in the following phase of a clinical trial: a) phase-1 clinical trials b) phase-2 clinical trials c) phase-3 clinical trials d) phase-4 clinical trials*	10 (18.5)	32 (59.26)	p<0.0001
8.	Select the correct (ADR and its causative drug) option: a)Phocomelia- Streptomycin b) Hemolytic anemia- Thalidomide c) HPA axis suppression - Ofloxacin d) Cleft lip- Phenytoin*	15 (27.7)	32 (59.26)	p<0.0001
9.	Select the correct (ADR and its causative drug) option : a) Yellowish discoloration of teeth- Isotretinoin b)Ebstein's cardiac anomaly- Warfarin c) Neural tube defects- Valproic acid* d)depressed nose, hand defects- Lithium	18 (33.3)	40 (74.1)	p<0.0001
10.	Regarding classification of ADR, the correct option is: a) Type A is predictable, dose related b) Type B is Unpredictable, dose unrelated c) Both a) and b) are correct* d) None of the above	20 (37)	38 (70.4)	p<0.0001
11.	It is important to report ADRs leading to- a) Hospitalization b) congenital abnormality c) patient death d) All of the above*	22 (40.7)	35 (64.8)	p<0.0001

Table 2. Attitude of Pharmacovigilance& ADR reporting before & after educational intervention.

Q.no	K A P Items	Pre-KAP Score n(%)	Post-KAP Score n(%)	p- value
12.	Do you agree that ADR reporting system would benefit patient care? -Strongly agree*	26 (48.2)	42 (77.7)	p<0.0001
13.	Would you suspect ADRs when drug is administered in normal dose? -Strongly agree*	19 (35.2)	43 (79.6)	p<0.0001
14.	Reporting of all ADRs for a new drug is essential? -Strongly agree*	17 (31.5)	50 (92.6)	p<0.0001
15.	Do you agree reporting of adverse drug reaction is necessary? -Strongly agree*	15 (27.7)	46 (85.2)	p<0.0001
16.	Do agree Pharmacovigilance should be taught in detail to healthcare professionals? -Strongly agree*	25 (46.3)	40 (74.1)	p<0.0001

Table3. Perception of Pharmacovigilance and ADR reporting before & after educational intervention.

Q.no	K A P Items	Pre-KAP Scores n(%)	Pre-KAP Scores n(%)	p- value
17.	Is it important to know national, international centers for ADR monitoring? -Yes*	15 (27.7)	40 (74.1)	p<0.0001
18.	Communication of safety information between all health care professionals can minimize the risk of marketed medicines? -Yes*	17 (31.5)	42 (77.7)	p<0.0001
19.	Can ADR monitoring help to promote rational use of medicines? -Yes*	18 (33.3)	44 (81.5)	p<0.0001
20.	Would you like to read an article (online /newspaper/Magazine) about ADR's in future? -Yes*	20 (37)	38 (70.4)	p<0.0001

Correct Response*, P<0.001 (comparison between the pre- KAP and Post- KAP responses).

Figure 1: Mean KAP scores of responders- Overall level of knowledge and attitude among the participants. (n=52)




Table.4. Student Feedback regarding educational intervention on Pharmacovigilance and Adverse drug reactions reporting:

Students opinions:	Response	n (%)
1. Sought information about Objectives of Pharmacovigilance-	Yes*	40 (74.1)
2. Sought information about Vigiflow database for adverse drug reactions monitoring-	Yes*	42 (77.7)
3. Sought essential information required while reporting an ADR-	Yes*	48 (88.8)
4. Sought information about different types of ADR-	Yes*	45 (83.3)
5. The hand-outs before the lecture helped us to grasp the ADR monitoring and Pharmacovigilance concepts during lecture better-	Yes*	36 (66.6)
6. Handouts before every Pharmacology lecture class helps to absorb concepts better-	Yes*	43 (79.6)
7. The photo images showing some examples of (ADR's and its causative drug) during lecture were informative-	Yes*	38 (70.4)

n- Number of Yes*responses, (%) - percentage of responses

CONCLUSION: In conclusion, the results of the present study demonstrate that an educational intervention can increase awareness of pharmacovigilance, ADR's reporting among the medical students and inculcate in their future

clinical practice. The medical students would be made aware about benefit- risk ratio of safety of marketed medicines and importance of communication with various health care professionals in pharmacovigilance.

REFERENCES:

1. Von Laue NC, Schwappach DL, Koeck CM., The epidemiology of preventable adverse drug events: a review of literature., Wien KlinWochenschr., 2003;115(12):407-415.
2. Wu WK, Pantaleo N et al., Evaluation of outpatient adverse drug reactions leading to hospitalization., American Journal of Health System Pharmacy.,2003;60(3):253-259.
3. Lazarou J, Pomeranz BH, Corey PN., Incidence of adverse drug reactions in hospitalized patients: a meta-analysis of prospective studies., Journal of the American Medical Association., 1998;279(15):1200-1205.
4. Pirmohamed M, James S, Meakin S, et al., Adverse drug reactions as cause of admission to hospital: prospective analysis of 18,820 patients., British Medical Journal., 2004;329(7456):15-19.

5. Figueiras A, Tato F, Fontainas J, Gestal-Otero JJ et al., Influence of physicians' attitudes on reporting adverse drug events: a case control study., *Medical Care.*, 1999;37(8):809-814.
6. Williams D, Feely J., Underreporting of adverse drug reactions: attitudes of Irish doctors., *Irish Journal of Medical Science.*, 1999;168:257-261.
7. Perlik F, Slanar O, Smid M, Petracek J et al., Attitude of Czech physicians to adverse drug reaction reporting., *European Journal of Clinical Pharmacology.*, 2002;58:367- 369.
8. Belton KJ, Lewis SC, Payne S, Rawlins MD et al., Attitudinal survey of adverse drug reaction reporting by medical practitioners in the United Kingdom., *British Journal of Clinical Pharmacology.*, 1995;39:223- 226.
9. Hasford J, Goettler M, Munter KH, Muller-Oerlinghausen B et al., Physicians' knowledge and attitudes regarding the spontaneous reporting system for adverse drug reactions., *Journal of Clinical Epidemiology.*, 2002;55:945-950.
10. Herdeiro MT, Figueiras A, Polo' nia J, Gestal-Otero JJ et al., Physicians' attitudes and adverse drug reaction reporting: a case control study in Portugal., *Drug Saf.*, 2005;28:825-833.
11. Hepler CD., Clinical pharmacy, pharmaceutical care, and the quality of drug therapy., *Pharmacotherapy.*, 2004;24:1491-8.
12. V.Lokesh Reddy., Assessment of Knowledge, Attitude and Perception of Pharmacovigilance and Adverse Drug Reaction (ADR) Reporting among the Pharmacy Students in South India., *IOSR Journal of Pharmacy and Biological Sciences.*, 2014;9(2):34-43.
13. Radhakrishnan R, Vidyasagar S, Varma DM., An Educational Intervention to assess Knowledge Attitude Practice of pharmacovigilance among Health care professionals in an Indian tertiary care teaching hospital., *Int.J. PharmTech Res.*, 2011;3(2):678-92.
14. Bagewadi Harish.G, Venkatadri.T.V, Nayaka Swapna. R., Impact of educational intervention on knowledge, attitude and perceptions of pharmacovigilance and adverse drug reactions in fifth term medical students and their feedback., *Int. J.of Res. in Pharmacology & Pharmacotherapeutics.*, 2015;4(2):164-174.
15. Cosentino M, Leoni O, Banfi F, Lecchini S, Frigo G., Attitudes to adverse drug reaction reporting by medical practitioners in a Northern Italian district., *Pharmacol Res.*, 1997;35:85-88.
16. Figueiras A, Herdeiro MT, Polonia J, Gestal-Otero JJ., An educational intervention to improve physician reporting of adverse drug reactions: a cluster-randomized controlled trial., *JAMA.*, 2006;296:1086-1093.
17. Suveges LG, Gesy KF, Wallace SM, Blackburn JL, Appel WC., Adverse drug reaction reporting part II: evaluation of the Saskatchewan pilot project for a regional reporting program in Canada., *Drug Information Journal.*, 1995; 29:581-589.
18. Scolt HD, Thacher – Renshaw A, Rosenbaum SE, Waters WJ, Green M, Andrews LG et al., Physician reporting of adverse drug reactions: results of the Rhode Island adverse drug reaction reporting project., *JAMA.*, 1990;263:1785-1788.
19. Chatterjee S, Lyle N, Ghosh S., A survey of the knowledge, attitude and practice of adverse drug reaction reporting by clinicians in eastern India., *Drug Saf.*, 2006;29:641-642.
20. Li Q, Zhang SM, Chen HT, Fang SP, Yu X, Liu D, Shi LY, Zeng FD., Awareness and attitudes of healthcare professionals in Wuhan, China to the reporting of adverse drug reactions., *Chin Med J.*, 2004;117:856-861.

Conflict of Interest Reported: Nil;

Source of Funding: None Reported